


Organic Carbon, Nitrogen, and Phosphorus Cycles and the composition of marine phytoplankton

1. C, N, P cycles
2. Major biochemicals in marine phytoplankton
3. Introduction to dissolved organic carbon

Major Carbon Reservoirs



Major Carbon Reservoirs

Atmosphere 750 GT C

Land Plants 900 GT C

Soils 1800 GT C

Carbonates 60,000,000 GT C


Kerogen 20,000,000 GT C

DIC 31,000 GTC


Marine Plants 2 GT C

Ocean 700 GT C


Sediments 150 GT C




Major Carbon Reservoirs



Particulate organic carbon (POC) and Dissolved organic carbon (DOC)



From Peltzer and Hayward, 1996



THE NITROGEN CYCLE




Figure by MIT OCW.

GLOBAL NITROGEN FIXATION (GT N/YR)

Industrial	0.05
Fossil Fuel Combustion	0.02
Lightning	0.01
Agriculture	0.09
Forests, etc.	0.05
Ocean	0.04

Figure by MIT OCW.

THE NITROGEN CYCLE (Tg N/year)




Figure by MIT OCW.

Marine Nitrogen Inventory (Excluding N₂) [2.3e7 Tg N]






Figure by MIT OCW.

Dissolved Nitrogen (μM) at HOTS



Preservoirs and Cycling in the Ocean



The pre-anthropogenic marine Phosphorus Cycle. see text for details on fluxes.

Figure by MIT OCW.

Elemental composition
and major biochemicals
of marine and terrestrial organisms
and the Redfield Ratio

Redfield Ratio

$$\text{C:N:P} = 106:16:1$$

Proteins	4:1:0
Carbohydrates	C only ?
Nucleic Acids	10:4:1
Lipids	C only

Proximate analysis of algal cells

Chlorophyceae (green algae)	Protein	Carbohydrate	Lipid	Ash
<i>Tetraselmis maculata</i>	72	21	7	(24)
<i>Dunaliella salina</i>	58	33	10	(8)
Chrysophyceae (golden brown algae)				
<i>Monochrysis lutheri</i>	53	34	13	(6)
<i>Syracosphaera carterae</i>	70	23	7	(37)
Bacillariophyceae (brown algae, diatoms)				
<i>Chaetoceros sp.</i>	68	13	16	(28)
<i>Skeletonema costatum</i>	58	33	10	(39)
<i>Coscinodiscus sp.</i>	74	16	10	(57)
<i>Phaeodactylum tricornerutum</i>	49	36	14	(8)
Dynophyceae (dinoflagellates)				
<i>Amphidinium carteri</i>	35	38	23	(14)
<i>Exuriella sp.</i>	37	44	20	(8)
Average	57	29	13	

Proteins

proteins serve as enzymes
and structural components of cells




Figure by MIT OCW.





Figure by MIT OCW.


amino acid



polymers of amino acids
C/N = 4, 80% of all nitrogen in marine algae
>50% of the carbon



PROTEIN ANALYSES

proteins can either be analyzed whole or after hydrolysis




Protein analyses by amino acid composition


Gel Electrophoresis of whole proteins



Amino Acid Composition of Marine Algae from Cowey and Corner (1966)




Amino Acid Composition of Zooplankton



Carbohydrates

CH_2O , no N or P. simple sugars = monosaccharides
2D and 3D polymers = polysaccharides
function as energy storage and cell structural components

Picture removed due to copyright restrictions.



Polysaccharide Analysis




Figure by MIT OCW.

Lipids

Distribution

ubiquitous. Many organisms have unique or characteristic suites of lipids, making them excellent tracers for sources of organic matter. Some lipids are also stable over 10^6 - 10^9 yr, and are therefore useful for paleoenvironmental reconstruction. Lipid mixtures in sediments and rocks are often extremely complex.

Function

Membrane components, energy storage, pigments, cell regulation

Chemical Structure

Polymers with C-C bonds, which are therefore very stable.
Lipids have the general formula C_nH_{2n} , normally with N,O,S,P
Lipids can be linear chains or cyclize to form rings, making them structurally diverse.

RH

ROH


RCOOH

RHNH₂

Lipid biosynthesis occurs via two pathways


1) polyketide

polymerization of acetate (CH_3COOH). products usually have an even number of carbons



2) isoprenoid

polymerization of isoprene a five carbon HC. Products have 10, 15, 20....carbons








Figure removed due to copyright restrictions.
Source Unknown.

Isoprenoid Biomarkers




α-terpinene

Monoterpenes C₁₀




phytol

Diterpenes C₂₀



Cholesterol


Triterpenes C₃₀



β-carotene

Tetraterpenes C₄₀

Terrestrial Plant organic components



Lignin is a complex structural polymer of hydroxycinnamyl alcohols that are linked together in a number of different linkages and patterns. Lignin and cellulose make up a large percentage of the carbon in woody and nonwoody tissues of higher plants.

Figure by MIT OCW.

DIN vs DIP, and TDN vs TDP at station ALOHA




Figure by MIT OCW.

Nitrogen versus phosphorus correlations for samples collected in the upper 0-400 m of the water column at Sta. ALOHA during the period October 1988 to December 1997. The bottom line is for nitrate plus nitrite (N+N) versus soluble reactive phosphorus (SRP) concentrations and the top line is for total dissolved nitrogen (TDN) versus total dissolved phosphorus (TDP). Model II linear regression analyses: $N+N (\mu M) = 14.62 [14.58 \text{ to } 14.66] SRP (\mu M) - 1.08 [-1.10 \text{ to } -1.06]$, $r = 0.996$, $n = 3299$ and $TDN (\mu M) = 14.57 [14.45 \text{ to } 14.69] TDP (\mu M) + 1.50 [1.44 \text{ to } 1.56]$, $r = 0.981$, $n = 2060$. Values in brackets indicate the 95% confidence intervals of the respective slope and intercept values.

DON and DOP at station ALOHA, HOTS






Figure by MIT OCW.

N/P ratios at station ALOHA, HOTS




Nitrogen-to-phosphorus (N:P) ratios versus water depth for samples collected at Sta. ALOHA during the period Oct. 1988 to Dec. 1997. (Left) Molar N:P ratios for dissolved inorganic pools calculated as nitrate plus nitrite (N + N): soluble reactive phosphorus (SRP). (Center) Molar N:P ratios for the "corrected" total dissolved matter pools. (Right) Molar N:P ratios for total dissolved matter pools, including both inorganic and organic compounds, calculated as total dissolved nitrogen (TDN) : total dissolved phosphorus (TDP). As a point for reference, the vertical dashed line in each graph is the Redfield-Ketchum-Richards molar ratio 16N:1P.

Major Carbon Reservoirs



TOC in the N. Pacific



Global inventory of DOC

DOC measured by high temperature catalytic oxidation (HTCO) method.

Global inventory was measured by International JGOFS program (680 GT C).

Depth pretty much the same everywhere, with high surface water values (40-80 $\mu\text{M C}$) and low deep water values (40 $\mu\text{M C}$).

Deep sea concentrations are nearly constant.






Figure by MIT OCW.


Radiocarbon Age of Dissolved Organic Carbon



DO¹⁴C and DI¹⁴C in the Atlantic Ocean



DO¹⁴C and DI¹⁴C in the Atlantic and Pacific Ocean



Radiocarbon measurements show that the average age of deep sea DOC is > 5000 yr. If the global inventory is 700 GT C, then the annual flux of C to maintain the system at steady state can be calculated as:


$$700 \text{ GT} / 5000 \text{ yr} = 0.14 \text{ GT C yr}^{-1}$$

The annual input of DOC from rivers is $0.2\text{-}0.4$ GT C yr^{-1} ,
While annual marine primary production is $60\text{-}75$ GT Cyr^{-1}

Either source adds enough C per year to maintain the system.

So what is the source of oceanic DOC?

Origin of DOC from stable carbon isotopes



A comparison of the stable isotope ratio measurements for DOC uv and suspended POC for the North Atlantic (Hydros-6) and the NCP (Eve-1) site.

Wrap up

The elemental composition (C, N, P, S) of marine and terrestrial organisms is a reflection of their biochemical composition at the molecular level.

Marine organisms (in terms of their carbon) are made of proteins > carbohydrates > lipids > nucleic acids. In terms of N, 70-80% is in proteins. The remaining is distributed between nucleic acids, carbohydrates, pigments, etc. The proportions of biochemicals are variable and reflect both environmental factors And physiological status.

N and P are limiting nutrients in the euphotic zone. Most of the N and P in the euphotic zone occur as DON and DOP. It is not known why these reservoirs of organic nutrients exist. Is the ocean N or P limited???

DOC is the largest reservoir of organic carbon in seawater. >98% of organic carbon in the ocean is DOC. It has high concentrations in the euphotic zone indicating net production or at least input. Values at depth are low (about half surface water values) and nearly constant.

Radiocarbon values of DOC are highly depleted, consistent with an average residence time of 5000-6000 years. Several ocean mixing cycles!!! No one knows why, but surely this impacts organic nutrients.