

## ABSTRACT ROOT SYSTEMS

YOUR NAME HERE

18.099 - 18.06 CI.

Due on Monday, May 10 in class.

*Write a paper proving the statements formulated below. Add your own examples, asides and discussions whenever needed.*

Let  $V$  be a Euclidean space, that is, a finite dimensional real linear space with a symmetric positive definite inner product  $\langle, \rangle$ .

Recall the definition of a reflection in  $V$  ([1]):

**Definition 1.** A reflection in  $V$  with respect to a vector  $\alpha \in V$  is defined by the formula:

$$s_\alpha(x) = x - \frac{2\langle x, \alpha \rangle}{\langle \alpha, \alpha \rangle} \alpha.$$

**Definition 2.** An abstract root system in  $V$  is a finite set  $\Delta$  of nonzero elements of  $V$  such that

- (1)  $\Delta$  spans  $V$ ;
- (2) for all  $\alpha \in \Delta$ , the reflections

$$s_\alpha(\beta) = \beta - \frac{2\langle \beta, \alpha \rangle}{\langle \alpha, \alpha \rangle} \alpha$$

map the set  $\Delta$  to itself;

- (3) the number  $\frac{2\langle \beta, \alpha \rangle}{\langle \alpha, \alpha \rangle}$  is an integer for any  $\alpha, \beta \in \Delta$ .

A root is an element of  $\Delta$ .

**Remark 3.** This definition may seem weird to you, but wait till you see the examples. A golden rule in mathematics is that if you can define an object for which there are enough, but not too many nontrivial examples, it must have important implications. If a nice classification of such objects is available, it is a discovery. This indeed is the case with the abstract root systems. They arise naturally in the theory of semisimple Lie algebras, and play an important role in a wide range of problems of algebra, representation theory and mathematical physics. At the same time, they can be defined and largely dealt with by means of linear algebra. This will be our goal.

---

Date: July 18, 2004.

**Example 4.** Let  $V$  be the following subspace of  $\mathbb{R}^{n+1}$ ,  $n \geq 1$ :

$$V = \left\{ \sum_{i=1}^{n+1} a_i e_i, \text{ with } \sum_{i=1}^{n+1} a_i = 0 \right\},$$

where  $\{e_i\}_{i=1}^{n+1}$  is an orthonormal basis in  $\mathbb{R}^{n+1}$ , and all  $a_i \in \mathbb{R}$ . Check that the set  $\Delta = \{e_i - e_j, i \neq j\}$ , is an abstract root system. This is the root system of type  $A_n$ . Describe geometrically (sketch) the set  $\Delta$  for  $n = 1$  and  $n = 2$ .

**Example 5.** Let  $V$  be the space  $\mathbb{R}^n$ ,  $n \geq 2$  with an orthonormal basis  $\{e_i\}_{i=1}^n$ . Check that the set

$$\Delta = \{\pm e_i \pm e_j, i \neq j\} \cup \{\pm e_i\}$$

is an abstract root system. This is the root system of type  $B_n$ . Describe geometrically the set  $\Delta$  for  $n = 2$ .

**Definition 6.** An abstract root system is reducible if it can be represented as a disjoint union of two abstract root systems  $\Delta = \Delta' \cup \Delta''$ , and each element of  $\Delta'$  is orthogonal to each element of  $\Delta''$ . We say that  $\Delta$  is irreducible if it admits no such decomposition.

**Example 7.** Check that the root systems of types  $A_2$  and  $B_2$  are irreducible. Check that a union of two root systems of type  $A_1$  is a reducible root system in  $V = \mathbb{R}^2$ , and sketch it.

We would like to classify the abstract root systems in any given dimension, but to do this we need to be precise about what it means for two root systems to be geometrically equivalent - we don't want to distinguish between a root system and the one obtained from it by a simple process like rotation or dilation. With this in mind, say that two root systems  $\Delta_1$  and  $\Delta_2$  are *isomorphic* if there is a linear automorphism  $f : V \rightarrow V$  taking  $\Delta_1$  to  $\Delta_2$ , such that for any roots  $\alpha$  and  $\beta$  in  $\Delta_1$ ,  $\frac{2\langle \beta, \alpha \rangle}{\langle \alpha, \alpha \rangle} = \frac{2\langle f(\beta), f(\alpha) \rangle}{\langle f(\alpha), f(\alpha) \rangle}$ . We will classify the abstract root systems "up to isomorphism", that is, treating isomorphic root systems as being the same.

To get an idea of how to make this classification, we start by proving some elementary properties.

**Theorem 8.** Let  $\Delta$  be an abstract root system in  $V$ .

- (1) If  $\alpha \in \Delta$ , then  $-\alpha \in \Delta$ .
- (2) If  $\alpha \in \Delta$  and  $\frac{1}{2}\alpha$  is not in  $\Delta$ , then the only possible elements of  $\Delta \cup \{0\}$  proportional to  $\alpha$  are  $\pm\alpha$ ,  $\pm 2\alpha$  and  $0$ .
- (3) If  $\alpha$  is in  $\Delta$  and  $\beta \in \Delta \cup \{0\}$ , then

$$\frac{2\langle \beta, \alpha \rangle}{\langle \alpha, \alpha \rangle} = 0, \pm 1, \pm 2, \pm 3, \text{ or } \pm 4,$$

and  $\pm 4$  can occur only if  $\beta = \pm 2\alpha$ .

Hint: use the Cauchy-Schwarz inequality in  $V$  ([1]) to prove (3).

**Example 9.** Compute the numbers  $n(\alpha, \beta) := \frac{2\langle\beta, \alpha\rangle}{\langle\alpha, \alpha\rangle}$  in two-dimensional examples of types  $A_2$ ,  $B_2$  and  $A_1 \oplus A_1$  considered in Example 7. Does this exhaust the possibilities predicted by the theorem? Can you guess any of the missing two-dimensional root systems?

Here is another statement to help in finding the root systems:

**Theorem 10.** Let  $\Delta$  be an abstract root system in  $V$ .

- (1) If  $\alpha$  and  $\beta$  are in  $\Delta$ , and  $\langle\alpha, \beta\rangle > 0$ , then  $\alpha - \beta$  is a root or 0. If  $\langle\alpha, \beta\rangle < 0$ , then  $\alpha + \beta$  is a root or 0.
- (2) If  $\alpha \in \Delta$  and  $\beta \in \Delta \cup \{0\}$ , then the set of elements of  $\Delta \cup \{0\}$  of the form  $\beta + n\alpha$ ,  $n \in \mathbb{Z}$ , contains all and only such elements with  $-p \leq n \leq q$ , for some  $p \geq 0$  and  $q \geq 0$  such that  $p - q = \frac{2\langle\beta, \alpha\rangle}{\langle\alpha, \alpha\rangle}$ .

Hint: To prove (2), assume there is a gap in the set of elements of  $\Delta \cup \{0\}$  of the form  $\beta + n\alpha$  and use (1) to get a contradiction.

What is the maximal number of roots contained in a set  $\beta + n\alpha$ , where  $\alpha \in \Delta$  and  $\beta \in \Delta \cup \{0\}$ ? (Use both theorems above).

Now use the Euclidean geometry. Recall that with the standard inner product in  $\mathbb{R}^n$ , the number  $\langle\alpha, \alpha\rangle = \|\alpha\|^2$  is the square of the length of the vector, and  $n(\alpha, \beta)$  can be written as

$$n(\alpha, \beta) = 2 \frac{\|\beta\|}{\|\alpha\|} \cos(\phi),$$

where  $\phi$  is the angle between  $\alpha$  and  $\beta$ .

Then we have

$$n(\alpha, \beta) \cdot n(\beta, \alpha) = 4 \cos^2(\phi).$$

Apply Theorem 8 to list the possibilities for angles  $\phi$  and relative lengths between two nonproportional elements of an abstract root system.

Using the above classification, construct examples of abstract root systems in  $V = \mathbb{R}^2$ . Remark: The one that contains two vectors with the angle between them  $\phi = \pi/6$  is called  $G_2$ .

#### REFERENCES

- [1] Your classmate, *Reflections in a Euclidean Space*, preprint, MIT, 2004.