

18.700 Problem Set 9

Due in class Tuesday December 3; late work will not be accepted. Your work on graded problem sets should be written entirely on your own, although you may consult others before writing.

1. (8 points) Suppose V is a real or complex inner product space. A linear map $S \in \mathcal{L}(V)$ is called *skew-adjoint* if $S^* = -S$. Suppose V is complex and finite-dimensional, and S is skew-adjoint. Show that the eigenvalues of S are all purely imaginary (that is, real multiples of i) and that there is an orthogonal direct sum decomposition

$$V = \bigoplus_{\lambda \in \mathbb{R}} V_{i\lambda}.$$

2. (16 points) Suppose V is an n -dimensional real vector space, and S is a skew-adjoint linear transformation of V .

- Show that Sv is orthogonal to v for every $v \in V$.
- Show that every eigenvalue of S^2 is a real number less than or equal to zero.
- Suppose (still assuming S is skew-adjoint) that $S^2 = -I$ (the negative of the identity operator on V). Show that we can make V into a *complex* inner product space, by defining scalar multiplication as

$$(a + bi)v = av + Sv$$

and the complex inner product as

$$\langle v, w \rangle_{\mathbb{C}} = \langle v, w \rangle - i\langle Sv, w \rangle.$$

What is the dimension of V as a complex vector space?

- Show that there is an orthonormal basis of V in which the diagonal blocks of the matrix of S are all either (0) or

$$\begin{pmatrix} 0 & -\lambda \\ \lambda & 0 \end{pmatrix}$$

with $\lambda > 0$. (In (d), we are **not** assuming that $S^2 = -I$.)

3. (5 points) Give an example of a square complex matrix A with the property that A has exactly three distinct eigenvalues, but A is *not* diagonalizable. (For full credit, you should *prove* that your matrix has the two required properties.)

MIT OpenCourseWare
<http://ocw.mit.edu>

18.700 Linear Algebra
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.